

CUMBU NAPIER HYBRID GRASS : YIELD, QUALITY AND SOIL FERTILITY STATUS AS INFLUENCED BY DIFFERENT NUTRIENT SOURCES

K. SATHIYA BAMA*

Department of Agronomy
Tamil Nadu Agricultural University,
Coimbatore (Tamil Nadu), India

*(e-mail : kssoilscience@gmail.com)

(Received : 10 October 2017; Accepted : 17 November 2017)

SUMMARY

To study the impact of different nutrient sources (organics, inorganics INM) on yield and quality of important perennial fodder, cumbu napier hybrid grass CO (CN)₄ an experiment was carried out. Three-year data of the experiment revealed that the basal application of FYM @ 105 t/ha (N equivalent basis)+2 kg *Azospirillum*+2 kg phosphobacteria recorded the highest green fodder yield of 360.7 t/ha/year. The crude protein content got increased in the FYM applied treatment. The higher organic carbon content of 1.28 per cent from the initial carbon status of 0.62 per cent recorded in the FYM treatment. The poultry manure (PM) applied plot recorded higher available N content of 185 kg/ha from the initial soil available N content of 165 kg/ha. The higher available K content of 578 kg/ha was recorded in the same treatment. The higher uptake of N, P, K, Ca, Mg, S, Fe, Mn, Zn and Cu was recorded in the FYM applied plots. The PM applied plot recorded the highest net returns of Rs. 2, 70,000/ha/year and B : C ratio of 4.3/ha.

Key words : Cumbu napier hybrid grass, farm yard manure, poultry manure, fodder yield, quality, nutrient uptake, soil nutrients, economics

Cumbu napier grass grows faster and produces more herbage. Nitrogen is an important agronomic input for bajra napier production. Historically, inorganic source has been a major N source. Now-a-days because of rising costs of natural gas which is used in N fertilizer synthesis, fuel for fertilizer transport, the organic manures also represent one of the sources of N which can be used in forages. The organic manures greatly enhanced the soil microbial population and produced more microbial biomass than inorganic which directly stimulated the activity of microorganism (Shanthi *et al.*, 2012). Manure can be a valuable source of nutrients for grass. It is a “complete” fertilizer, containing varying amounts of all the major and minor plant nutrients. Manure applications may also improve soil physical properties such as water infiltration rate, aggregate size and stability, pore size and crust strength. These improvements in turn translate into better soil quality and improved productivity. Besides the differences in N concentration, different N sources may not be equally effective when applied to fodder crops. A number of factors, including soil type, rate and method of application, forage management and environmental

conditions can impact the effectiveness of different fertilizer sources to provide N to crops (Mubarak *et al.*, 2010).

The required amount of nutrients for napier grass may be supplied through both inorganic and organic for grass. But due to the farmers' demand on organic way of grass fodder production, the research work is started with organic sources alone and compared with integrated nutrient management and inorganics.

The recently developed cumbu napier hybrid grass variety CO (CN)₄ has the greatest potential to grow under various soil fertility and nutrient conditions. In the present research work, the easily available organic sources such as FYM and poultry manures are tried as N source. FYM is a well recognized and considered as balanced bulky organic manure which increases the moisture retention of the soil and helps in dissolution of nutrients. The organics were applied along with bio-fertilizers (*Azospirillum*). The beneficial effect of *Azospirillum* on plant is associated not only with the process of N fixation and improved nutrition of plants but also with the synthesis of complex biologically active compounds such as

pyridoxine, biotin, gibberellins and other compounds which accelerate plant growth under favourable environmental conditions. Among the various nutrient management practices, the current issue of identifying suitable organic source to cumbu napier grass will be highly useful to the 30 organic forage growers. Of the major nutrient elements, nitrogen has special significance in increasing green biomass yield and its quality. Nitrogen fertilization has been reported to improve not only the yield but also the crude protein content. With the above points in mind, this study was attempted with different locally available organic sources such as FYM and poultry manure on N equivalent basis and compared with INM and inorganics to get sustained optimum yield. Bama (2014) reported the higher carbon sequestration potential of the cumbu napier hybrid grass type. Results of the field experiment revealed that the application of spentwash @ 50 kilo litre per ha at full dose with recommended dose of nitrogen and phosphorus increased the quality parameters, nutrient parameters and green fodder yield over recommended dose of fertilizers (Latha *et al.*, 2015).

MATERIALS AND METHODS

The experiment was carried out at the Department of Forage Crops, Tamil Nadu Agricultural University, Coimbatore, Tamil Nadu to study the impact of different nutrient sources (organics, inorganics INM) on yield and quality of important perennial fodder cumbu napier hybrid grass CO(CN)₄ during the years 2009 to 2012. The nutrient sources viz., farm yard manure (FYM) on N equivalent basis (105 t/ha) with 2 kg *Azospirillum*+2 kg phosphobacteria (S₁), poultry manure (44 t/ha) with 2 kg *Azospirillum*+2 kg phosphobacteria (S₂), INM (Inorganics+25 t FYM/ha/year) (S₃) and inorganics alone (75 : 50 : 50 kg N, P and K/ha/year as basal and 75 kg N/ha/year after each harvest, totally seven cuts/year) (S₄) 525 kg N/ha were tried in this experiment. The FYM and poultry manure applied on N equivalent basis and for inorganics treatment, NPK recommendation was 75 : 50 : 50 kg/ha. Each time N was top dressed @ 75 kg/ha and for INM (inorganics+25 t of FYM/ha) and inorganics. Totally seven cuts were harvested per annum and total N requirement @ 525 kg/ha/year was applied. The soil analysis report before the commencement of experiment revealed that the soil was slightly alkaline (7.8) and free from excessive salts (0.12 dS/m). It was medium in organic carbon (0.65%) and low in

available N (165 kg/ha), medium in available P (15 kg/ha) and high in available K (510 kg/ha). The green fodder yield was recorded as such in the field. The dry fodder yield was recorded by keeping the samples in electric oven at 65°C. The plant samples were analyzed at cutting stage for N, P, K, S, Ca, Mg, S and micronutrients by acid digestion technique (Piper, 1966). The fodder quality parameters viz., crude protein, crude fibre and crude fat contents were estimated by the procedure outlined by Van Soest and Moore (1965). The data were analyzed statistically by using simple RBD.

RESULTS AND DISCUSSION

Fodder Yield

The first year data of this experiment showed that among the different sources of nutrients tried to improve the green fodder yield of cumbu napier hybrid grass, the FYM applied on N equivalent basis (105 t FYM/ha/year) recorded the highest green fodder yield of 341.3 t/ha/year followed by poultry manure applied on N equivalent basis (44 t PM/ha/year) which recorded 314 t/ha/year (Table 1). Though the INM (282.4 t/ha/year) recorded numerically higher yield than inorganics, it was statistically on par with inorganics alone (275.2 t/ha/year). During second and third years' also, same trend of results was observed. The pooled analysis of three years' data showed that the application of FYM @ 105 t/ha/year recorded 360 t/ha/year which was 25 percent increase over inorganics (fig1). The higher dry fodder yield was recorded in FYM (79.18 t/ha/year) and the lowest was recorded in the inorganics alone (63.85 t/ha/year). These findings are in close conformity to those of and Kumar and Sharma (2002) in fodder sorghum. The higher green fodder yield in the organics applied plot might be attributed to better supply of nutrients by conducive physical environment leading to better root activity and higher nutrient absorption, which resulted in better plant growth and superior yield (Mishra *et al.*, (2008), Thakur *et al.* (2011). Golada *et al.* (2012) reported that application of FYM significantly increased the green fodder yield of pearl millet.

Quality Parameters

Fodder quality

The important fodder quality parameters viz., crude protein, crude fibre and crude fat contents were

TABLE 1
Effect of source of nutrients on yield and quality yield cumbu napier hybrid grass

Treatment	Green fodder yield(t/ha/yr)				Dry fodder yield (t/ha/yr)				Quality (%)		
	I year	II year	III year	Mean	I year	II year	III year	Mean	Crude protein	Crude fibre	Crude fat
S ₁ -FYM	373.5	341.3	365.2	360.7	86.3	79.18	84.73	83.4	8.8	31.9	2.3
S ₂ -PM	350.2	314.0	342.1	335.4	80.9	72.84	79.37	77.7	8.3	31.5	2.2
S ₃ -INM	325.7	282.4	314.3	307.5	75.2	65.52	72.92	71.2	8.1	32.1	2.1
S ₄ -Inorganics	256.9	275.2	269.4	267.2	59.3	63.85	62.50	61.9	7.6	32.5	2.1
C. D. (P=0.05)	20.1	26.8	21.3		5.1	6.8	6.3		0.4	NS	NS

NS-Not Significant.




Fig. 1. Influence of nutrient sources on green fodder yield (t/ha) analyzed. Among the treatments, the FYM applied plot recorded higher crude protein content (8.8%). Bama *et al.* (2013) reported that poultry manure application @18.8 t/ha+2 kg *Azospirillum*+2 kg phosphobacteria recorded 167.6 t green fodder yield/ha/year. There was no significant difference observed in the crude fibre and crude fat analytical data (Table 1).

Nutrient uptake

The data on the uptake of nitrogen indicated that there was significant increase in FYM applied plot of 1215 kg/ha which was followed by PM applied (1074 kg/ha) and INM (974 kg/ha) (Table 2). The higher P uptake of 518 kg/ha was recorded in the FYM plot which was on par with PM applied plot (502 kg/ha). The lowest P uptake was recorded in the inorganic plot (332 kg/ha). The potassium uptake recorded during first year indicated that there was significant increase in FYM applied plot of 1476 kg/ha recorded which was on par with PM applied (1383 kg/ha) followed by INM (1286 kg/ha). The lowest K uptake was recorded in inorganic plot of 1014 kg/ha (Fig. 2). Forage crops require high amount of N and P (Latha *et al.*, 2013). The increase in uptake of nutrients in the organic applied plots may be due to slow release of nutrients by the organics might have met the fodder crop requirement (Suganya, 2008).

Highest Ca uptake of 967 kg/ha was recorded

in the FYM plot which was followed by INM applied plot (842 kg/ha) and PM plot (825 kg/ha). The lowest Ca uptake recorded in inorganic alone plot (664 kg/ha). The uptake of magnesium indicated that the FYM applied plot recorded 328 kg/ha which was on par with PM applied plot (324 kg/ha) followed by INM plot (271 kg/ha). The lowest Mg uptake was recorded in inorganic plot of 213 kg/ha. The sulphur uptake recorded during first year indicated that there was significant increase of S uptake in FYM applied plot of 95 kg/ha which was on par with PM applied (88 kg/ha) followed by INM (82 kg/ha). The lowest N uptake was recorded in inorganic plot of 65 kg/ha. The increased secondary nutrients such as Ca, Mg and S uptake might be due to higher biomass production by the organic application as well as higher nutrient content in the organics.




Fig. 2. Influence of nutrient sources on major nutrient uptake (kg/ha)

Highest Fe uptake of 176 kg/ha was recorded in the FYM plot which was on par with PM applied plot (165 kg/ha) followed by INM plot (153 kg/ha). The lowest Fe uptake was recorded in inorganic plot (121 kg/ha). The uptake of manganese indicated that the FYM applied plot recorded 23.0 kg/ha which was on par with PM applied plot (21.5 kg/ha) and followed by INM plot (20.0 kg/ha). The lowest uptake was recorded in inorganic plot of 15.8 kg/ha. Higher Zn uptake of 15.9 kg/ha was recorded in the FYM plot

TABLE 2
Effect of source of nutrients on nutrient uptake of Cumbu Napier hybrid grass

Treatment	Nutrient uptake (kg/ha/year)									
	N	P	K	Ca	Mg	S	Fe	Zn	Mn	Cu
S ₁ -FYM	1215	518	1476	967	328	95	176	15.9	23.0	4.83
S ₂ -PM	1074	502	1383	825	324	88	165	14.9	21.5	4.53
S ₃ -INM	975	451	1286	842	271	82	153	13.8	20.0	4.21
S ₄ -Inorganics	721	332	1014	664	213	65	121	10.9	15.8	3.32
C. D. (P=0.05)	102	56	152	96	35	9	19	1.8	2.0	NS

NS-Not Significant.

which was on par with PM applied plot (14.9 kg/ha) followed by INM plot (13.8 kg/ha). The lowest Zn uptake was recorded in inorganics plot (10.9 kg/ha). The same trend of results was observed in second and third year also. Non-significant copper uptake was observed. Organics had significant influence on the micro-nutrients compared to recommended dose of fertilizer. This might be due to high micro-nutrient content with FYM application. This is in line with the findings of and Madhumitadas *et al.* (2010). The increased nutrient uptake by the poultry manure alone applied plot in the multicut sorghum was reported by Bama *et al.* (2013).

Soil fertility

In the cumbu napier hybrid grass, there was a drastic improvement in the organic carbon status of the soil by the application of organic manures (Table 3). The FYM applied on N equivalent basis recorded higher organic carbon content of 1.28 per cent, followed by poultry manure applied treatment (0.91%), from the initial carbon status of 0.62 per cent. The increase in organic carbon content in the manorial treatment was attributed to direct addition of organic manure in the soil which stimulated the growth and activity of microorganisms and also due to better root growth resulting in the higher production of biomass, crop stubbles and residues (Yilmaz and Alagoz, 2010;

Singh *et al.*, 2011; Moharana *et al.*, 2012). The subsequent decomposition of these materials might have resulted in the enhanced carbon content of soil. These results are in agreement with the findings of Singh *et al.* (2008) and Kumar *et al.* (2008). Further the addition of FYM might have created environment conducive for formation of humic acid, which ultimately resulted in an increase in the organic carbon content of the soil of the field. The analytical data of the available nitrogen status of the soil showed that the poultry manure applied plot recorded higher available N content of 185 kg/ha followed by FYM applied treatment of 180 kg/ha compared to rest of the treatments. The increase in available N might be attributed to the enhanced multiplication of microbes by the incorporation of manures for the conversion of organically bound N to inorganic form and release of N bearing organic compound during FYM decomposition. The favourable soil conditions under organic manure application might have facilitated the mineralization of soil N leading to build-up of higher available N (Vipin Kumar and Singh, 2010). The higher available P content of 20.2 kg/ha was observed in the poultry manure applied plot which was followed by FYM applied plot (19.8 kg/ha). The increased availability of P with organics could be ascribed to their solubilizing effect on the native soil P and consequent contribution of the P as solubilized to labile pool. Organic matter may also reduce the fixation of

TABLE 3
Effect of source of nutrients on soil nutrient status of Cumbu napier hybrid grass grown soil

Treatments	SOC (%)	Major nutrients (kg/ha)			Secondary nutrients cmol p(+)/kg		DTPA extractable micronutrients (ppm)			
		N	P	K	Ex.Ca	Ex.Mg	Fe	Zn	Mn	Cu
S ₁	1.28	180	19.8	578	17.5	7.2	10.2	6.8	7.2	7.2
S ₂	0.91	185	20.2	561	15.2	6.8	8.7	5.9	6.7	6.8
S ₃	0.80	160	19.5	553	14.3	6.2	7.9	5.1	6.2	6.2
S ₄	0.70	153	18.5	532	13.5	5.7	7.8	4.5	5.7	5.7
C. D. (P=0.05)	0.10	17	0.3	16	1.8	0.8	0.7	0.5	0.8	0.8

phosphate by providing protective cover and reduce the phosphate fixing capacity and increase the available P in soil of the field.

Higher available K content of 578 kg/ha was recorded in the FYM applied treatment. The higher exchangeable Ca content was recorded in the FYM plot (17.5 cmol (p+)/kg) which was followed by PM plot (15.2 cmol (p+)/kg). The INM (14.3 cmol (p+)/kg) and inorganics (13.5 cmol (p+)/kg) recorded comparable values. Higher exchangeable Mg was recorded in FYM plot (7.2 cmol (p+)/kg) which was followed by 6.8 cmol (p+)/kg. The DTPA extractable micro-nutrient results revealed that higher Fe content was recorded in FYM applied plot (10.2 ppm) followed by INM (7.9 ppm) and inorganics (7.8 ppm). The same trend of results was observed in Zn, Mn and Cu contents.

Soil nutrient balance sheet

The major nutrients available in the soil after completing three year period indicated that there was a drastic depletion of almost all nutrients in the soil (Table 4). The nutrient uptake pattern of cumbu napier hybrid grass indicated that N, P and K content was removed heavily from the soil. The secondary and micronutrients were also heavily removed from the soil. If that crop is recommended widely the fertilizer schedule has to be restudied. Especially the potassium was highly mined from the soil. Hence, the K study related to pools of K, sources and doses of K had to be taken care. Bama (2016) also reported same results with lucerne crop.

Effect of sources of nutrients on economics of fodder cultivation

Data pertaining to influence of inorganic and

organic sources of nutrients on economics of fodder cultivation are presented in Table 5. The results showed that the PM applied plot recorded the highest net returns (Rs. 2,70,000) and B : C ratio (4.3) per ha followed by INM (Rs. 2,46,915) and PM (Rs. 2,28,500). But the FYM application recorded lower B/C ratio of 2.5. Apart from economics according to Bama and Babu (2016), forages particularly grass type fodder contributed to carbon sequestration and most part of the long time carbon storage from roots i. e. below ground portion. This type of fodder crop can saturate carbon level quickly wherever the climate change mitigation is essential. Among the various forage crops, cumbu napier hybrid grass removed higher carbon removal by above ground biomass and in the below ground biomass. Among the sources, farm yard manure. This information adds an another important point to this article.

CONCLUSION

The experiment taught that the basal application of FYM @105 t/ha+2 kg *Azospirillum*+2 kg phosphobacteria may be recommended for highest green fodder yield and crude protein yield and soil fertility. However, from economic point of view, if the farmer wants to adopt organic farming practice, the PM 44 t/ha+2 kg *Azospirillum*+2 kg phosphobacteria can be recommended. However, the negative nutrient balance of almost all nutrients revealed that the grass was a voracious feeder of nutrients. Hence these nutrients have to be recommended with required dose to grass type fodder crops. If grass type cumbu napier hybrid grass is recommended for cultivation, the fertilizer schedule has to be researched.

TABLE 4
Soil nutrient balance sheet for Cumbu Napier Hybrid Grass

Treatment	Initial (a)	Applied (b)	Uptake (c)	Post-harvest (d)	Actual gain (d-a)	N balance (a+b)-(c+d)
Nitrogen						
S ₁	165	525	1174	180	-15	-664
S ₂	165	525	1032	185	-20	-527
S ₃	165	525	923	160	5	-393
S ₄	165	525	753	153	12	-216
Phosphorus						
S ₁	15	210	500	19.8	-4.8	-294.8
S ₂	15	176	482	20.2	-5.2	-311.2
S ₃	15	50	427	19.5	-4.5	-381.5
S ₄	15	50	347	18.5	-3.5	-300.5
Potassium						
S ₁	510	525	1426	578	-68	-969
S ₂	510	176	1329	561	-51	-1204
S ₃	510	50	1218	553	-43	-1211
S ₄	510	50	1058	532	-22	-1030

TABLE 5
Effect of source of nutrients on monitory benefits of different fodder crops

Treatment	Cumbu napier			
	CC	GR	NR	B/C
S ₁ -FYM	145000	373500	228500	2.5
S ₂ -PM	80200	350200	270000	4.3
S ₃ -INM	78785	325700	246915	4.1
S ₄ -Inorganics	53785	256900	203115	4.7

CC- Cost of cultivation (Rs), GR- Gross return (Rs),NR- Net return (Rs),B/C-Benefit cost ratio.

REFERENCES

- Bama, S. K. 2014 : Prediction of carbon sequestration potential of forage system. *J. Ecobiol.* **33** : 169-175.
- Bama, S. K. 2016 : Enshot of different nutrient sources on fodder yield, quality and soil fertility status of lucerne grown soil. *Forage Res.*, **41** : 222-227.
- Bama, S. K., and C. Babu. 2016 : Perennial forages as a tool for sequestering atmospheric carbon by best management practices for better soil quality and environmental safety. *Forage Res.*, **42** : 149-157
- Bama, S. K., K. Velayudham, C. Babu, K. Iyanar, and A. Kalamani. 2013 : Enshot of different nutrient sources on fodder yield, quality and soil fertility status of multicut fodder sorghum grown soil. *Forage Res.*, **38** : 207-212. <http://forageresearch.in>.
- Golada, S. L., G. L. Sharma, A. Verma, and H. K. Jain. 2012 : Effect of FYM, nitrogen and Azospirillum on yield, economics and soil nutrient status of forage pearl millet. *Madras Agric. J.* **99** : 308-310.
- Kumar, B., R. K. Gupta, and A. L. Bhandari. 2008 : Soil fertility changes after long-term application of organic manures and crop residues under rice-wheat system. *J. Indian Soc. Soil Sci.* **56** : 80-85.
- Kumar, S., and B. L. Sharma. 2002 : Effect of FYM, nitrogen and Azospirillum inoculation on yield and quality of fodder sorghum. *Forage Res.*, **28** : 165-168.
- Latha, P., P. Thangavel, K. Velayudham, and K. Arulmozhiselvan. 2013 : Effect of distillery industry byproducts on nutrient and enhanced yield of cumbu napier hybrid grass. *J. Indust. Pollut. Control*, **29** : 73-79.
- Madhumitadas, H., R. B. Chakraborty, Singandhupe, S. D. Muduli, and A. Kumar. 2010 : Utilization of distillery wastewater for improving production in under productive paddy grown area in India. *J. Sci. Indust. Res.* **69** : 560-563.
- Mishra, B., A. Sharma, S. K. Singh, J. Prasad, and B. P. Singh. 2008 : Influence of continuous application of amendments to maize-wheat cropping system on dynamics of microbial biomass in Alfisol of Jharkhand. *J. Indian Soc. Soil Sci.* **56** : 71-75.
- Moharana, P. C., B.M. Sharma, D. R. Biswas, B. S. Dwivedi, and R. V. Singh. 2012 : Long-term effect of nutrient management on soil fertility and soil organic carbon pools under a 6-year-old pearl millet-wheat cropping system in an Inceptisol of sub-tropical India. *Field Crops Res.*, **136** : 32-41.
- Mubarak, A. R., A. Eshraga, M. Gali, G. Asma Mohamed, D. Steffens, and H. Abdelkarim Awadelkarim. 2010 : Nitrogen mineralization from five manures as influenced by chemical composition and soil type. *Commun. Soil Sci. and Plant Analysis*, **41** : 1903-1920.
- Piper, C. S. 1966 : *Soil and Plant analysis*. Hans Publishers, Bombay.
- Shanthi, G., P. E. Somasundaram, S. Muthukrishnan and S.Thiyageshwari. 2012 : Influence of organic sources of nutrients on soil fertility and quality parameters of turmeric. *Madras Agric. J.* **99** : 553-555.
- Singh, P., R. Kumar, and S. Pal. 2008 : Integrated nutrient management in rice-wheat cropping system for sustainable productivity. *J. Indian Soc. Soil Sci.* **56** : 205-208.
- Singh, R. N., Surendra Singh, S. S. Prasad. V. K. Singh, and Pramod Kumar. 2011 : Effect of integrated nutrient management on soil fertility, nutrient uptake and yield of rice-pea cropping system on an upland acid soil of Jharkhand. *J. Indian Soc. Soil Sci.* **59** : 158-163.
- Suganya, K. 2008 : Recycling options of one time application of distillery spentwash to maize and groundnut and its impact on soil fertility and groundwater quality. Ph. D. (Env. Sci.) thesis, Tamil Nadu Agricultural University, Coimbatore.
- Thakur, R., S. D. Sawarkar, U. K. Vaishya, and M. Singh. 2011 : Impact of continuous use of inorganic fertilizers and organic manure on soil properties and productivity under soybean-wheat intensive cropping of Vertisol. *J. Indian Soc. Soil Sci.* **59** : 74-81.
- Van Soest, P. J., and L. A. Moore. 1965 : New chemical methods for analysis of forages for the purpose of predicting nutritive value. *Proc. IXth Intenat. Grassl. Congr.* **9** : 783-789.
- Kumar, V. and A. K. Singh. 2010 : Long-term effect of green manuring and farm yard manure on yield and soil fertility status in rice-wheat cropping system. *J. Indian Soc. Soil Sci.* **58** : 409-412.
- Yilmaz, and Alagoz. 2010 : Effects of short-term amendments of farm yard manure on some soil properties in the Mediterranean region-Turkey. *J. Food, Agric. and Environ.*, **8** : 859-862.