

## GROWTH, PRODUCTIVITY AND PROFITABILITY OF FODDER PEARL MILLET AS INFLUENCED BY VARIOUS NUTRIENT MANAGEMENT PRACTICES

RAKESH KUMAR, HARDEV RAM\*, PRAVEEN B. R.<sup>1</sup>, CHETHAN BABU R. T.<sup>2</sup>, SUPRIYA AND BHARTI DEVI<sup>3</sup>

ICAR-National Dairy Research Institute, Karnal-132 001 (Haryana), India

<sup>1</sup>Tea Research Association, North Bengal Regional R & D Centre, Nagrakata, Jalpaiguri-735 225 (West Bengal), India

<sup>2</sup>Teaching Associate, Department of Agronomy, S. V. Agricultural College, Tirupati-517502, India

<sup>3</sup>G.S.G.D Girls Agriculture College, Padampur-335041, India

\*(e-mail: [devagron@gmail.com](mailto:devagron@gmail.com))

(Received: 2 September 2024; Accepted: 25 September 2024)

### SUMMARY

A field experiment was conducted during *kharif* season of 2019-20 at Research Farm, Agronomy Section, ICAR-National Dairy Research Institute, Karnal (Haryana). The experiment was laid out in Randomized Complete Block Design with eight treatments, *i.e.*, T<sub>1</sub> (Absolute control); T<sub>2</sub> (100% RDF); T<sub>3</sub> (100% RDF + Cow urine foliar spray); T<sub>4</sub> (100% RDF + PGPR); T<sub>5</sub> (100% RDF + PGPR + Cow urine foliar spray); T<sub>6</sub> (75% RDF + Cow urine foliar spray); T<sub>7</sub> (75% RDF + PGPR) and T<sub>8</sub> (75% RDF + PGPR + Cow urine foliar spray) with three replications. Study indicated that the growth and yield parameters of fodder pearl millet significantly affected with different nutrient management practices and found plant height (178.3 and 307.5 cm), leaf length (86.5 and 110.4 cm), no. of leaves/plant (11.4 and 15.3), no. of tillers/plant (11.3 and 13.1), leaf width (4.3 and 5.1 cm) and stem girth (2.2 and 3.7 cm) at 40 DAS and harvest respectively, while, yield of green (54.58 t/ha) and dry fodder yield (11.17 t/ha) at harvest significantly increase up to application of 100% RDF+PGPR and they further increase with application of 100% RDF+PGPR+CU, but doesn't reach up to significance level over 100% RDF+PGPR. However, both were remains statistically at par to each and found significantly higher than other treatments. Maximum net return was recorded with T<sub>4</sub> followed by T<sub>5</sub>, T<sub>2</sub> and T<sub>7</sub> treatment. However, maximum B:C ratio was found with T<sub>4</sub>, followed by T<sub>7</sub> and T<sub>5</sub> treatments. Which, will further strengthen and sustain crop productivity and profitability.

**Key words:** Cow urine, Fodder, Growth, Pearl millet, Yield

Livestock production is important pillar of Indian agriculture and About 20.5 million people rely on livestock for a living, and animals contributed 16 percent of small farm households' income, compared to 14 percent for all rural households. India has the highest number of livestock animals (536.76 million) in the world. The population of major livestock animals *viz.*, buffalo, cattle, sheep and goat in India is 109.85, 193.46, 74.26 and 148.88 million, respectively (Kumar *et al.*, 2022). Fodder demand increases for ever increasing population of livestock and its essential component for livestock production as it can cope up the cost of feeding because feeding accounts for 65 to 70 percent of the entire cost of livestock farming. The scarcity of green fodder is severe, and India alone faces a net deficiency 35.6, 10.95 and 44 percentage of green fodder, dry fodder and concentrate feeds

(Kumar *et al.*, 2023), respectively, that influence production levels as well as health of animals, which ultimately affect returns from livestock sector (Surve *et al.*, 2011). The availability of good quality green fodder throughout the year to livestock is the major concern to developing a sustainable livestock farming (Chaudhary *et al.*, 2016). Among the different fodder crops pearl millet (*Pennisetum glaucum*) is the gifted crops to tropical and sub-tropical regions that provide fodder, stover and food to millions of poor farmer and their livestock's. It has potential to grow in low fertility soils (Ali, 2010), higher growth rate, water use efficiency, tillering potential, and heat tolerance (Jukanti *et al.*, 2016), higher degree of tolerance to drought (Ibrahim *et. al.*, 2014).

Green revolution scenario shows production and productivity of cereals largely increased through

intensive agronomic practices, high yielding variety and Indiscriminate use of higher rate of chemical fertilizers with little or without use of organic source of nutrients to plant, that create adverse effect on soil properties *viz.*, inadequacy in one or more nutrients and deterioration of soil fertility which leads to stagnating or even declining crop productivity (Kumar *et al.*, 2023). Judicious use of organic and inorganic nutrients sources may sustain and enhance the crop quality and productivity. Cow urine and PGPR are great and vital sources of nutrients for plants among many organic sources. The cow urine contains; N, P, K, S, Na, Fe, Mn, Si, carbolic acid, hormones and phytohormones (Saunders, 1982). PGPR is a liquid consortium of microorganisms, which contains a wide range of genera *viz.*, *Pseudomonas Azospirillum*, *Bacillus*, *Serratia* and *Azotobacter* (Bashan *et al.*, 2004) that actively colonize around roots of plant and enhances plant growth and yield (Wu *et al.*, 2005) due to their ability to produce various phytohormones (Auxins, gibberellins, cytokinin and ethylene), organic acids, siderophores, biologically fixation of atmospheric nitrogen, increase solubility and availability of insoluble inorganic soil phosphate, oxidation of sulphur, extra cellular production of antibiotics, increases root permeability leads to enhance essential plant nutrients uptake (Enebak and Carey, 2000 and Pal *et al.*, 1999). Perceiving above facts the present study was proposed to finds out a suitable combination of nutrients sources to enhance productivity and profitability of fodder pearl millet.

## MATERIALS AND METHODS

### Description of experimental site

The experiment was carried out during *khariif*, 2019 at Agronomy Research Farm, ICAR-NDRI, Karnal (Haryana) located at 29°45' North latitude and 76°58' East longitude and at an altitude of 245 m above mean sea level. The area's climate is semi-arid, with an average annual rainfall of 707mm, 70-80 percent of it falling between July and September. During this investigation, the average minimum and maximum temperatures were 20.49°C and 34.54°C, respectively. The soil texture at the experimental site was clay loam (Piper, 1942), electrical conductivity (EC) 0.37 dS/m (Jackson, 1967), organic carbon (OC) 0.49 percent (Walkley and Black's, 1934), available nitrogen 215 kg/ha (Subbiah and Asija, 1956), available phosphorus (P) 24.70 kg/ha (Olsen *et al.*, 1954), and available potassium 285 kg/ha (Jackson, 1967).

### Treatment details and inputs application

The experiment was laid out in Randomized Complete Block Design with eight treatments, *i.e.*, T<sub>1</sub> (Absolute control), T<sub>2</sub> (100% RDF), T<sub>3</sub> (100% RDF + Cow urine foliar spray), T<sub>4</sub> (100% RDF + PGPR), T<sub>5</sub> (100% RDF + PGPR + Cow urine foliar spray), T<sub>6</sub> (75% RDF + Cow urine foliar spray), T<sub>7</sub> (75% RDF + PGPR) and T<sub>8</sub> (75% RDF + PGPR + Cow urine foliar spray) with three replications. The land preparation involved one deep ploughing with disc plough followed by two cross harrowing with disc harrow and planking. The recommended dose of fertilizers (80:30:30 kg/ha, N: P<sub>2</sub>O<sub>5</sub>: K<sub>2</sub>O, respectively) were applied according to treatments. The half of N and full doses of P and K were applied before final harrowing and remaining half dose of nitrogen was top-dressed in two split doses as per the treatment. The PGPR (100 ml/10 kg seeds) liquid culture was diluted in one litres of water, and applied on seeds. Thereafter, inoculated seeds were dried in shade for 60-90 minutes, after drying seeds was manually sown. Nutrifed variety of fodder pearl millet was sown using 10 kg seed per hectare with maintaining row to row spacing 30 cm and plant to plant spacing 10 cm. Other package of practices was followed as per standard for cultivation of fodder pearl millet. The 10% cow urine was applied as foliar spray in early morning hours, when the dew has been evaporated at 30 and 45 DAS as per treatments.

### Pre and post-harvest observations

Five plants were chosen at random and tagged in the net plot area to capture morphological parameters across treatments. Plant height, leaf width and length of middle leaf, number of leaves/plants, number of tillers/plants and stem girth were measured at 40 days after sowing and harvest. while, leaf to stem ratio and green fodder yield were recorded at harvest. The crop was harvested manually at 50 % flowering stage. Net plot area was harvested separately from each plot, weighed as kg/plot and then converted into tonnes/ha for estimation of final fresh green fodder yield.

$$\text{Leaf to stem ratio} = \frac{\text{Mean fresh leaves weight of five tagged plants}}{\text{Mean fresh stem weight of five tagged plants}}$$

### Statistical analysis

All of the data was analysed using analysis of variance (Gomez and Gomez 1984) at a 5% level of significance ( $P < 0.05$ ). Simple Pearson's correlation coefficient was computed by using mean values of different growth and yield parameters.

## RESULTS AND DISCUSSION

### Morphological traits and leaf to stem ratio

Growth parameters *i.e.*, plant height, number of leaves and tiller per plant, leaf width, leaf length and stem girth are trustworthy indicator of growth and development of any plant, especially for fodder crops, which directly or indirectly represents final yield of fodder crops. Study indicated (Table 1, 2 and 3) that plant growth parameters of fodder pearl millet were significantly influenced with different nutrient

management practices and recorded significantly higher in all the treatments at 40 days after sowing and harvest over absolute control. Plant height (178.3 and 307.5 cm), leaf length (86.5 and 110.4 cm), no. of leaves/plant (11.4 and 15.3), no. of tillers/plant (11.3 and 13.1), leaf width (4.3 and 5.1 cm) and stem girth (2.2 and 3.7 cm) at 40 DAS and harvest respectively, while leaf to stem ratio (0.44) at harvest significantly increase up to application of 100% RDF+PGPR ( $T_4$ ) and they further increase with application of 100% RDF+PGPR+CU ( $T_5$ ) but doesn't reach up to significance level over 100% RDF+PGPR ( $T_4$ ). However, both were remains statistically at par to each and found significantly higher than other treatments. Nutrients supplied with 100% RDF+PGPR improved plant height by 11.25, 13.66 and 55.85%; leaf length by 6.91, 7.81 and 27.88%; leaf width by 6.77, 10.79, and 36.9%; no. of tillers/plant by 11.90, 14.49 and 61.22%; no. of leaves/plant by 13.02, 15.32 and 53.33% and stem girth by 12.21, 15.83 and 69.00%

TABLE 1  
Effects of nutrient management practices on morphological characteristics of fodder pearl millet

Treatments	Plant height (cm)		Leaf length (cm)		No. of leaves/plant	
	40 DAS	Harvest	40 DAS	Harvest	40 DAS	Harvest
$T_1$ : Absolute control	107.4	197.3	59.2	86.3	7.3	10.0
$T_2$ : 100% RDF	155.5	270.5	77.5	102.4	10.0	13.3
$T_3$ : 100% RDF+CU	158.5	276.4	78.0	103.2	10.2	13.5
$T_4$ : 100% RDF+PGPR	178.3	307.5	86.5	110.4	11.4	15.3
$T_5$ : 100% RDF+PGPR+CU	181.0	308.7	86.7	111.4	11.5	15.7
$T_6$ : 75% RDF+CU	127.1	228.9	68.0	94.2	8.5	11.4
$T_7$ : 75% RDF+PGPR	147.2	261.9	76.6	101.6	9.8	12.9
$T_8$ : 75% RDF+PGPR+CU	150.7	267.0	76.9	102.1	9.9	13.1
S. Em( $\pm$ )	6.42	9.91	2.67	2.39	0.36	0.46
C. D. ( $P=0.05$ )	19.47	30.05	8.11	7.25	1.10	1.40

Note: CU: Cow urine; DAS: Day after sowing; RDF: Recommended dose of fertiliser and PGPR: Plant growth promoting rhizobacteria

TABLE 2  
Effects of nutrient management practices on leaf width, number of tillers and stem girth of fodder pearl millet

Treatments	Leaf width (cm)		No. of tillers/plant		Stem girth (cm)	
	40 DAS	Harvest	40 DAS	Harvest	40 DAS	Harvest
$T_1$ : Absolute control	3.0	3.7	6.8	8.1	1.2	2.1
$T_2$ : 100% RDF	3.9	4.6	9.8	11.5	1.8	3.2
$T_3$ : 100% RDF+CU	3.9	4.7	9.9	11.7	1.9	3.3
$T_4$ : 100% RDF+PGPR	4.3	5.1	11.3	13.1	2.2	3.7
$T_5$ : 100% RDF+PGPR+CU	4.4	5.3	11.4	13.4	2.3	3.8
$T_6$ : 75% RDF+CU	3.4	4.1	8.5	9.9	1.5	2.6
$T_7$ : 75% RDF+PGPR	3.8	4.5	9.6	11.1	1.7	3.0
$T_8$ : 75% RDF+PGPR+CU	3.8	4.5	9.7	11.4	1.8	3.1
S. Em( $\pm$ )	0.09	0.10	0.34	0.39	0.07	0.11
C. D. ( $P=0.05$ )	0.29	0.31	1.04	1.18	0.21	0.34

over 100% RDF+CU, 100% RDF and absolute control at harvest, respectively.

According to data, growth parameters increase due to optimum and balanced supply of essential nutrients to plant throughout the growth period by RDF, PGPR and cow urine. Optimum and balanced supply of phosphorus in early stage of plant from fertiliser, and PGPR that responsible for early and well establishments of rooting system of plant that acquire more water and nutrients from different strata of soil and supply to plant for profuse and healthy growth. Optimum availability of nitrogen in soil solution, increase uptake, utilization and assimilation leads to increase chlorophyll synthesis in plant because nitrogen is an integral and important component of porphyrin ring for formation of chlorophyll. Higher chlorophyll molecules per unit area of leaf, intercept more solar radiation and it convert into organic compound through photosynthesis mechanism with the help of water, carbon dioxide (CO<sub>2</sub>). Photosynthates that synthesized during photosynthesis process accumulate mostly in leaves and stem with the help of phloem, resulted increases cell size, cell division, expansion and differentiation. Cell division increase number of cells leads to produce a greater number of nodes, which responsible to emerge a greater number of tillers, internodes and leaves per plants. Higher accumulation of photosynthate in cells, which responsible to increase cell size leads to increase leaf length and width; and internodes length and diameters. Higher internodes length and diameters attributed to increase plant height and stem girth (Iqbal *et al.*, 2017). Potassium protects to plant from diseases, pests and play important role in plant to maintain water status. The similar results also reported by Chattha *et al.* (2017) and Bhakar *et al.* (2021).

In addition to RDF, PGPR is another source of nutrient to plant, that actively colonize around plant roots (Wu *et al.*, 2005) and increase native phosphorus availability by solubilization of insoluble inorganic soil phosphate into soluble form, supplement additional nitrogen by biological fixation of atmospheric nitrogen throughout the growth period (Enebak and Carey, 2000), release ammonia into soil subsequence increase soil nitrate (Abbasi *et al.*, 2011), increase K<sup>+</sup>/Na<sup>+</sup> ratio in plant by suppressing Na<sup>+</sup> uptake (Ahmad *et al.*, 2014) and increase metal nutrients availability in soil to plant by organic acid production (Cakmakci *et al.*, 2007) these all mechanism recharge soil fertility status as well as increase essential plant nutrients in available form to plant use. At same time cow urine additionally supplement N, K, S, P and micronutrients to plant

foliage in latter stage of crop for quick recovery (Sadhukhan *et al.*, 2018). In another way PGPR synthesis phytohormones (Auxin, gibberellin and cytokinin) (Kumar *et al.*, 2014); secrete ACC-deaminase that leads to decrease ethylene levels in plant (Ahmad *et al.*, 2014); and produce HCN secondary metabolites that suppress germination of pathogen spores (Kumar *et al.*, 2014); control bacterial diseases through extra cellular production of antibiotic (Pal *et al.*, 1999); increase nutrient and water uptake due to increases in root permeability (Glick, 1995); eliminate deleterious rhizobacteria from the rhizosphere by niche exclusion (Weller, 1988); and cow urine contains hormones and enzymes (Kishore *et al.*, 2015); cow urine foliar spray control fungal and bacterial diseases (Devakumar *et al.*, 2014) these all stimulatory mechanism increase physiological process of plant attributed to increase growth by decreasing stress and deleterious effect from different factor's on plant.

### Fodder yield

Study indicated (Table 3) that green and dry fodder yield of fodder pearl millet were significantly influenced with different nutrient management practices. Yield of green fodder (54.58 t/ha) and dry fodder (11.17 t/ha) at harvest significantly increase up to application of 100% RDF+PGPR (T<sub>4</sub>) and they further increase with application of 100% RDF+PGPR+CU (T<sub>5</sub>) but doesn't reach up to significance level over 100% RDF+PGPR (T<sub>4</sub>). However, both were remains statistically at par to each and found significantly higher than other treatments. Nutrients supplied with 100% RDF+PGPR, increase green fodder yield by 5.65, 6.50 and 59.37% over 100% RDF+CU, 100% RDF and absolute control at harvest, respectively. Green fodder yield per unit area

TABLE 3  
Effects of nutrient management practices on fodder yield and leaf to stem ratio of fodder pearl millet

Treatments	Green fodder yield (t/ha)	Dry fodder yield (t/ha)	Leaf to stem ratio
T <sub>1</sub> : Absolute control	34.25	5.56	0.37
T <sub>2</sub> : 100% RDF	51.25	9.89	0.41
T <sub>3</sub> : 100% RDF+CU	51.66	10.02	0.42
T <sub>4</sub> : 100% RDF+PGPR	54.58	11.17	0.44
T <sub>5</sub> : 100% RDF+PGPR+CU	54.83	11.33	0.44
T <sub>6</sub> : 75% RDF+CU	47.41	8.23	0.39
T <sub>7</sub> : 75% RDF+PGPR	50.85	9.40	0.41
T <sub>8</sub> : 75% RDF+PGPR+CU	51.16	9.61	0.41
S. Em(±)	8.39	1.90	0.003
C. D. (P=0.05)	2.54	0.57	0.010

is out come from interaction of cultivar potential, agronomic management practices and environmental factors that are directly and indirectly contributes in final yield. The green fodder yield is the prominent factor to decide efficacy of applied input and agronomic management practices. Balanced and regular supply of essential plant nutrients increases plant physiological processes leads to increase plant height; higher no. of leaves and tillers/plant; higher leaf length, width and stem girth responsible to increase final green fodder yield. Yield of dry fodder increases due to higher accumulation of photosynthates in plant that leads to increase dry matter content (Chattha *et al.*, 2017). The similar findings also reported by Chattha *et al.* (2017) and Bhakar *et al.* (2021).

### ECONOMICS

The practical serviceability's of any treatment is judged by the net returns and returns gained per rupee invested, *i.e.*, benefit cost ratio. So, with this objective, the economics of different treatments was worked out. Results depicted (Table 4) that the highest net returns found with application of 100% RDF+PGPR (Rs. 57124 ha<sup>-1</sup>) followed by 100% RDF+PGPR+CU (Rs. 56099 ha<sup>-1</sup>), 100% RDF (Rs. 52364 ha<sup>-1</sup>) and 75% RDF+PGPR (Rs. 52351 ha<sup>-1</sup>). However, maximum B:C ratio was found with application of 100% RDF+PGPR (2.31) followed by 75% RDF+PGPR (2.19), 100% RDF+PGPR+CU (2.15) and 100% RDF (2.14). Conjoint application of different source of nutrients to plants, found positively correlated and their higher efficacy gave higher fodder yield leads to achieve higher net return and returns gained per rupee invested. The similar findings also

TABLE 4  
Effects of nutrient management practices on economics of fodder pearl millet production

Treatments	Cost of cultivation (Rs./ha)	Gross return (Rs./ha)	Net return (Rs./ha)	B:C ratio
T <sub>1</sub> : Absolute control	20353	51375	31022	1.52
T <sub>2</sub> : 100% RDF	24511	76875	52364	2.14
T <sub>3</sub> : 100% RDF+CU	25911	77500	51589	1.99
T <sub>4</sub> : 100% RDF+PGPR	24751	81875	57124	2.31
T <sub>5</sub> : 100% RDF+PGPR+CU	26151	82250	56099	2.15
T <sub>6</sub> : 75% RDF+CU	25096	71125	46029	1.83
T <sub>7</sub> : 75% RDF+PGPR	23936	76288	52351	2.19
T <sub>8</sub> : 75% RDF+PGPR+CU	25336	76750	51414	2.03

reported by Chattha *et al.* (2017) and Bhakar *et al.* (2021).

### Correlation studies

The results showed on correlation (Table 5) indicates that the dry fodder yield ( $r=0.978$ ), height of plant ( $r=0.923$ ), length of leaf ( $r=0.935$ ), width of leaf ( $r=0.894$ ), no. of tillers/plant ( $r=0.932$ ), no. of leaves/plant ( $r=0.897$ ) and stem girth ( $r=0.906$ ) were strongly and positively correlated with the yield of green fodder. Similar findings also reported by Bhakar *et al.* (2021).

### CONCLUSION

Nutrient management is an important aspect, where intensive cropping system are dominates. The better nutrient management is pre-requisite to sustain the crop productivity and profitability. Based on the findings of present investigation, it was concluded that among different nitrogen management practices,

TABLE 5  
Correlation coefficient (r) between fodder yield and morphological traits of pearl millet at harvest.

Pearson	Correlations							
	GFY	DFY	PH	LL	LW	TP	LP	SG
GFY	1							
DFY	.978**	1						
PH	.923**	.982**	1					
LL	.935**	.986**	.998**	1				
LW	.894**	.965**	.989**	.988**	1			
TP	.932**	.986**	.997**	.998**	.993**	1		
LP	.897**	.968**	.995**	.994**	.995**	.996**	1	
SG	.906**	.974**	.997**	.995**	.997**	.997**	.998**	1

Note: GFY: Green fodder yield; DFY: Dry fodder yield; PH: Plant height; LL: Leaf length; LW: Leaf width; TP: Tillers per plant; LP: Leaves per plant; SG: Stem girth and \*\*=Correlation is significant at the 0.01 level.

combined application of 100% RDF + PGPR can successfully sustained crop productivity and profitability. Judicious use of inorganic and organic sources of nutrients sustains and enhance crop productivity and profitability and positive effect of integration was noticed on growth, yield as well as on economics of fodder pearl millet cultivation.

#### ACKNOWLEDGEMENTS

We sincerely acknowledge Indian Council of Agricultural Research for financial support and Director, ICAR-NDRI, Karnal for providing field staff, laboratory facilities and assistance in conducting this research.

#### REFERENCES

- Abbasi, M. K., S. Sharif, M. Kazmi, T. Sultan and M. Aslam, 2011: Isolation of plant growth promoting rhizobacteria from wheat rhizosphere and their effect on improving growth, yield and nutrient uptake of plants. *Plant Biosystems*, **145**(1): 159-168.
- Ahmad, M., Z. A. Zahir, M. Jamil, F. Nazli, M. Latif and M. F. Akhtar, 2014: Integrated use of plant growth promoting rhizobacteria, biogas slurry and chemical nitrogen for sustainable production of maize under salt-affected conditions. *Pakistan Journal of Botany*, **46**(1): 375-382.
- Ali, E. A., 2010: Grain yield and nitrogen use efficiency of pearl millet as affected by plant density, nitrogen rate and splitting in sandy soil. *American-Eurasian Journal of Agricultural and Environmental Sciences*, **7**: 327-335.
- Bashan, Y., G. Holguin and L. E. De-Bashan, 2004: Azospirillum-plant relationships: physiological, molecular, agricultural, and environmental advances 1997-2003. *Canadian Journal of Microbiology*, **50**(8): 521-577.
- Bhakar, A., M. Singh, S. Kumar, R. K. Meena, B. L. Meena, R. Kumar and V. K. Meena, 2021: Growth, productivity and profitability of fodder sorghum and cluster bean as influenced by mixed cropping and nutrient management. *Legume Research-An International Journal*, **44**(11): 1308-1314.
- Cakmakci, R., M. F. Donmez and U. Erdogan, 2007: The effect of plant growth promoting rhizobacteria on barley seedling growth, nutrient uptake, some soil properties, and bacterial counts. *Turkish Journal of Agriculture and Forestry*, **31**(3): 189-199.
- Chattha, M. B., A. Iqbal, M. U. Chattha, M. U. Hassan, I. Khan, I. Ashraf, M. Faisal and M. Usman, 2017: PGPR inoculated-seed increases the productivity of forage sorghum under fertilized conditions. *Journal of Basic and Applied Sciences*, **13**: 150-153.
- Chaudhary, D. P., A. Kumar, R. Kumar, A. Singode, G. Mukri, R. P. Sah, U. S. Tiwana and B. Kumar, 2016: Evaluation of normal and specialty corn for fodder yield and quality traits. *Range Management and Agroforestry*, **37**(1): 79-83.
- Devakumar, N., S. Subha, G. G. E. Rao and J. I. Khan, 2014: Studies on soil fertility, cow urine and panchagavya levels on growth and yield of maize. *Building Organic Bridges*, **2**: 627-630.
- Enebak, S. A. and W. A. Carey, 2000: Evidence for induced systemic protection to Fusarium rust in Loblolly pine by plant growth promoting rhizosphere. *Plant Disease*, **84**: 306-308.
- Glick, R. B., 1995: The enhancement of plant growth promotion by free-living bacteria. *Canadian Journal of Microbiology*, **41**: 109-117.
- Gomez, K. A. and A. A. Gomez, 1984: Statistical Procedures for Agricultural Research, *John Willey and Sons*, Singapore, pp. 680.
- Ibrahim, Y. M., A. E. Idris and M. A. E. Marhoum, 2014: Effect of Nitrogen Fertilizer on Irrigated Forage Pearl Millet (*Pennisetum Americanum* L.K. Shcum). *Universal Journal of Agricultural Research*, **2**(2): 56-60.
- Iqbal, A., M. A. Iqbal, A. Iqbal, Z. Aslam, M. Maqsood, Z. Ahmad, N. Akbar, H. Z. Khan, R. N. Abbas, R. D. Khan, G. Abbas and M. Faisal, 2017: Boosting forage yield and quality of maize (*Zea mays* L.) with multi-species bacterial inoculation in Pakistan. *International journal of experimental botany*, **86**: 84-88.
- Jackson, M. L., 1967: Soil Chemical Analysis. Prentice Hall of India Private Limited, New Delhi, pp. 111-203.
- Jukanti, A. K., C. L. Gowda, K. N. Rai, V. K. Manga and R. K. Bhatt, 2016: Crops that feed the world Pearl Millet (*Pennisetum glaucum* L.): an important source of food security, nutrition and health in the arid and semi-arid tropics. *Food Security*, **8**(2): 307-329.
- Kishore, S. V., L. R. Rao, B. Ramesh and A. K. Aditya, 2015: Indian cow urine distillation and therapeutic uses. *Mintage Journal of Pharmaceutical and Medical Sciences*, **4**(1): 1-5.
- Kumar, A., B. R. Maurya and R. Raghuvanshi, 2014: Isolation and characterization of PGPR and their effect on growth, yield and nutrient content in wheat (*Triticum aestivum* L.). *Biocatalysis and Agricultural Biotechnology*, **3**(4): 121-128.
- Kumar, R., H., Ram, R. K., Meena, S., Kumar, B., Kumar, Praveen, B. R. & P. S. Hindoriya, 2022: Nutrients content, uptake and soil biological properties as influenced by various nutrient management

- practices under fodder pearl millet cultivation. *Indian Journal of Ecology*, **49**(6): 2119-2124.
- Kumar, R., H., Ram, R. K., Meena, S., Kumar & V. K. Meena, 2023: Nutritional value and energy balance of pearl millet fodder as influenced by different nutrient management practices. *Indian J Dairy Sci*, **76**(3): 284-288.
- Kumar, R., H., Ram, R., Kumar, R. K., Meena, B. L., Meena & D. Kumar, 2023: Proximate composition and fibre fraction of pearl millet fodder as influenced by different nutrient management practices. *Indian Journal of Animal Research*, **57**(3): 334-339.
- Olsen, S. R., C. V. Cole, F. S. Watanabe and L. A. Dean, 1954: Estimation of available phosphorus in soils by extraction with sodium bicarbonate. USDA Circular 939, U.S. Government Printing Office, Washington D.C., pp. 19.
- Pal, K. K., R. Dey, D. M. Bhatt and S. M. Chauhan, 1999: Enhancement of groundnut growth and yield by plant growth promoting rhizobacteria. *International Arachis Newsletter*, **19**: 51-53.
- Piper, C. S., 1942: Soil and Plant Analysis: *A laboratory manual of methods for the examination of soils and the determination of the inorganic constituents of plants*. Hans, Bombay, pp. 152.
- Sadhukhan, R., J. S. Bohra and S. Choudhury, 2018: Effect of fertility levels and cow urine foliar spray on growth and yield of wheat. *International Journal of Current Microbiology and Applied Sciences*, **7**(3): 907-912.
- Saunders, W. H. M., 1982: Effects of cow urine and its major constituents on pasture properties. *New Zealand Journal of Agricultural Research*, **25**(1): 61-68.
- Subbiah, B. V. and G. L. Asija, 1956: A rapid procedure for determination of available nitrogen in soil. *Current Science*, **25**: 259-260.
- Surve, V., P. Patil and M. Arvadia, 2011: Forage production potential of sorghum (*Sorghum bicolor*), maize (*Zea mays*) and cowpea (*Vigna unguiculata*) under sole and intercropping systems. *Madras Agricultural Journal*, **98**(4): 372-374.
- Walkley, A. and I. A. Black, 1934: An examination of the Degtjareff method for determining soil organic matter, and a proposed modification of the chromic acid titration method. *Soil science*, **37**(1): 29-38.
- Weller, D. M., 1988: Biological control of soilborne pathogens in the rhizosphere with bacteria. *Annual Review of Phytopathology*, **26**: 379-407.
- Wu, S. C., Z. H. Cao, Z. G. Li, K. C. Cheung and M. H. Wong, 2005: Effects of biofertilizer containing N-fixer, P and K solubilizers and AM fungi on maize growth: a greenhouse trial. *Geoderma*, **125**(1-2): 155-166.