

PERFORMANCE OF FORAGE AND LEGUMES INTERCROPPING IN *JATROPHA CURCAS* PLANTATION IN SEMI-ARID REGION OF HARYANA

M. K. SINGH*, MAHAVIR PARSHAD, R. K. ARYA¹, PAWAN KUMAR² AND K. S. AHLAWAT

Department of Forestry,
CCS Haryana Agricultural University,
Hisar-125 004 (Haryana), India

*(e-mail : mks.hau.cssri@gmail.com)

(Received: 28 January, 2013, Accepted: 9 July, 2013)

SUMMARY

A field experiment was conducted to identify suitable crops for *Jatropha* based intercropping system during winter season of 2005-06 and 2006-07 at Department of Forestry Farm, CCS HAU, Hisar. The production potential of greengram (*Vigna radiata*). cv-MH-96-1, clusterbean (*Cyamopsis tetragonoloba*). cv-HG-563, mothbean (*Phaseolus aconitifolius*). cv-RMO-40 and pearl millet (*Pennisetum typhoides*). cv-HHB-67 were evaluated under 5m x 3m spacing of *Jatropha* planted in September 2003. *Jatropha* plantation had no significant effect on the growth and yield of test crops during the first year of experimentation except that mothbean, branching and yield was significantly reduced over control. However, the growth, yield and yield attributes of all the test crops were reduced significantly in *Jatropha* based intercropping system over control during the second year of experimentation. Two year old *Jatropha* produced negligible seed yield both during 2005-06 and 2006-07 due to frost injury in 2005-06 and excessive vegetative growth during 2006-07. Therefore, susceptibility of *Jatropha* to frost, requirement of irrigation for flowering and fruiting and poor seed yield has rendered it unsuitable for north India.

Key words : Intercropping, greengram, clusterbean, mothbean, pearl millet and *Jatropha curcas*

India is not self sufficient in petroleum production and nearly 75% of its requirement are being met through imports. Therefore, there is an urgent need for finding out alternate sources of energy which are renewable, safe and non-polluting. After rigorous study and research, oil extracted from different plant species was tested as fuel which could emerge as a strong bio-fuel with minimum effect on environment. All these characteristics were found in one species, called, *Jatropha*.

Jatropha curcas L (family Euphorbiace) is a multipurpose large shrub or small tree. It grows on well drained soil with good aeration and is well adapted to marginal soils with low nutrient content. It grows as a boundary fence or live hedge and can be used to reclaim eroded areas (Joker and Jepsen, 2003). Its leaves and stems are toxic to animals. So, it is not browsed, but after treatment, the seed or seed cake can be used as an

animal feed. Being rich in nitrogen, the seed cake is also an excellent source of plant nutrients (Makkar *et al.*, 2001).

Agroforestry is a modern tool to develop sustainable land use and to increase food production by growing woody species (trees, shrubs, bamboos etc) with agricultural crops and / or animals in some form of spatial arrangement or temporal sequence. Because these species co-exist crucial to determine the success of an agroforestry system. A survey of the available information reveals that most of the agroforestry species have negative allelopathic effects on food and fodder crops. Allelopathy and important ecological phenomenon play a significant role in diverse ecosystems. Allelochemicals-the chemicals involved in interplant interactions process diversity in terms of nature and structure. The phenomenon has been reported and agricultural systems are known to be allelopathic. In

¹Agriculture Development Officer, Bhuna.

²Department of Genetics and Plant Breeding.

³Department of Agronomy.

agroforestry system the role of allelopathy is especially important as it may lead to soil sickness and may be a casual factor for declining crop productivity reported by Batish *et al.*, (2011)

Keeping in view the importance of Jatropha oil, Jatropha plantation is being promoted by different agencies. It could be intercropped with other crops plants, however, meagre work has been done in relation to intercropping of food crops with Jatropha. The present study was, therefore, undertaken to find out the suitable summer crops which could be grown with *Jatropha curcas* in interspaces in semi-arid conditions.

MATERIALS AND METHODS

Field experiment was conducted during winter season of 2005-06 and 2006-07 at the Research Farm of Department of Forestry, CCS Haryana Agricultural University, Hisar located at 29°10'N latitude and 75°46'E longitude with an elevation of 215.2 m above the mean sea level. The climate of the experimental site is semi-arid with dry hot summer, cold winter and receives 452 mm average annual rainfall. Soil of the experimental field was sandy loam in texture, slightly saline in nature, low in nitrogen, and medium in phosphorus and rich in potassium. The field experiment consisted of *Jatropha curcas* planted in September, 2003 at 5m x 3m spacing intercropped with greengram (*Vigna radiata*). cv-MH-96-1, clusterbean (*Cyamopsis tetragonoloba*). cv-HG-563, mothbean (*Phaseolus aconitifolius*). cv-RMO-40 and pearl millet (*Pennisetum typhoides*). cv-HHB-67. The treatments were replicated five times in Randomized Block Design. In between the inter spaces of Jatropha plantation all the test crops were sown in middle of July with spacing of 45 x 30cm in both years. The recommended package of practices for the test crops were followed both in control and Jatropha. All growth, yield and yield attributes of test crops were recorded at the time of final harvest and analysed statistically. Plant height and branches/plant of Jatropha were recorded during the month of October and clusters/plant and fruits/plant were recorded in October and December months during both the years. Picking of the mature fruits was done at regular intervals from October to January. The Jatropha plants were cut back in March with the help of saw at above 45-60 cm above the ground due to killing of the above parts by severe frost in the first fortnight of January 2006. All the plants sprouted again in the month of April-May.

RESULTS AND DISCUSSION

The results of the present study revealed that Jatropha plantation had no adverse effect on growth, yield attributes and yield of greengram (Table 1), clusterbean (Table 2) and pearl millet (Table 4) during the first year of experimentation. However, the primary and secondary branches as well as grain and stover yield of mothbean (Table 3) were significantly reduced even during the first year of experimentation. It could be ascribed to more sensitivity of mothbean to shade and below ground interferences of Jatropha. Sharma (2003) has also reported 94 percent decrease in seed yield of mothbean compared to 71 and 72 percent seed yield decrease in mungbean and horsebean, respectively. Under 7 year old *Acacia tortilis* intercropping system compared to sole cropping. Due to increase in crown size and increased competition of roots for moisture and nutrients, the growth, yield attributes and grain as well as stover yield of all the test crops were significantly reduced in association with Jatropha over control during the second year of experimentation. Divya *et al* (2006) have also reported reduced plant height and grain yield of intercrops i.e. groundnuts, blackgram, cowpea, frenchbean, sunflower and gingelly under Jatropha plantation at different spacings. Rizvi *et al.*, (1999) have found that mimosine inhibited large number of physiological and biochemical parameters in *V. Mango* and *P. Aureus*. They found that mimosine inhibited seedling vigor, food mobilization efficiency, solubilisation of starch, breakdown of proteins and activity of amylase. The reduced amylase activity was at synthetic as well as catalytic level and it was mediated by gibberellic acid. They further reported that mimosine altered the hormonal balance of the seedlings leading to an inhibition in their growth. When *V. Mango* plants were grown in the soil having different amounts of leucaena leaves, nitrogenase activity of root nodules was inhibited.

Jatropha after two years of plantation (2005-06) produced negligible mean seed yield of 0.16 q/ha with no variation in seed yield between sole crop and intercrop (Table 5). The poor seed yield was due to severe damage to the fruits on account of frost condition (-3.5°C) in the first fortnight of January, 2006. Singh *et al* (2009) has also reported susceptibility of Jatropha to frost and its ability to sprout again in spring. The number of fruits/cluster (3.6) was also reduced markedly over the previous (2004-05) year (9.6 fruits/cluster) with negligible variation between intercropped Jatropha and sole Jatropha. It was

TABLE 1
Effect of Jatropha plantation on growth, yield and yield attributes of intercropped Greengram

Treatment	Primary branches/plant		Secondary branches/plant		No. of grains/pod		Test weight (1000-seeds) (g)		Grain yield (q/ha)		Stover yield (q/ha)		Harvest index (%)	
	05-06	06-07	05-06	06-07	05-06	06-07	05-06	06-07	05-06	06-07	05-06	06-07	05-06	06-07
year														
Greengram	3.10 (5.16)	3.24 (18.83)	4.79 (2.71)	4.96 (20.16)	24.96 (5.77)	25.64 (15.29)	10.09 (4.56)	10.20 (6.67)	11.86 (4.30)	12.37 (10.75)	27.28 (2.24)	28.32 (9.04)	30.02 (0.87)	30.39 (2.04)
Jatropha + Greengram	2.94	2.63	4.66	3.96	23.52	21.72	9.63	9.52	11.35	11.04	26.67	25.76	29.76	29.77
C. D. at 5%	NS	0.48	NS	0.08	NS	0.05	NS	0.27	NS	0.24	NS	0.58	NS	NS

TABLE 2
Effect of Jatropha plantation on growth, yield and yield attributes of intercropped Clusterbean

Treatment	Primary branches/plant		Secondary branches/plant		No. of grains/pod		Test weight (1000-seeds) (g)		Grain yield (q/ha)		Stover yield (q/ha)		Harvest index (%)	
	05-06	06-07	05-06	06-07	05-06	06-07	05-06	06-07	05-06	06-07	05-06	06-07	05-06	06-07
year														
Clusterbean	8.41 (2.97)	8.74 (13.50)	12.96 (3.63)	13.26 (9.80)	8.86 (1.24)	9.23 (5.85)	22.98 (2.09)	23.29 (8.59)	15.14 (1.45)	15.71 (17.76)	40.42 (1.48)	41.63 (13.60)	26.59 (0.79)	26.45 (0.23)
Jatropha + Clusterbean	8.16	7.56	12.49	11.96	8.75	8.69	22.50	21.29	14.92	12.92	39.82	35.97	26.38	26.39
C. D. at 5%	NS	0.66	NS	0.52	NS	0.27	NS	0.73	NS	0.28	NS	0.63	NS	NS

The data in parenthesis indicate % reduction over control.

TABLE 3
Effect of Jatropha plantation on growth, yield and yield attributes of intercropped Mothbean

Treatment	Primary branches/plant		Secondary branches/plant		No. of grains/pod		Test weight (1000-seeds) (g)		Grain yield (q/ha)		Stover yield (q/ha)		Harvest index (%)	
	05-06	06-07	05-06	06-07	05-06	06-07	05-06	06-07	05-06	06-07	05-06	06-07	05-06	06-07
Mothbean	7.64 (13.87)	7.88 (19.92)	5.26 (16.16)	5.52 (23.91)	5.59 (5.90)	5.71 (10.51)	27.51 (4.03)	28.52 (8.94)	5.96 (13.42)	6.07 (18.29)	14.60 (10.62)	14.18 (10.16)	30.96 (0.58)	30.87 (0.62)
Jatropha + Mothbean	6.58	6.31	4.41	4.20	5.26	5.11	26.40	25.97	5.16	4.96	13.05	12.74	30.78	30.68
C. D. at 5%	0.87	1.09	0.68	0.69	NS	0.53	NS	1.21	0.13	0.39	0.41	0.20	NS	NS

TABLE 4
Effect of Jatropha plantation on growth, yield and yield attributes of intercropped Pearl millet

Treatment	Primary branches/plant		Secondary branches/plant		No. of grains/pod		Test weight (1000-seeds) (g)		Grain yield (q/ha)		Stover yield (q/ha)		Harvest index (%)	
	05-06	06-07	05-06	06-07	05-06	06-07	05-06	06-07	05-06	06-07	05-06	06-07	05-06	06-07
Pearl millet	29.65 (6.37)	30.51 (19.01)	21.67 (5.45)	22.70 (17.36)	25.28 (5.46)	25.71 (11.63)	9.50 (3.79)	9.85 (9.24)	28.78 (6.81)	29.79 (14.94)	72.52 (5.69)	75.66 (14.26)	28.60 (0.03)	28.66 (0.07)
Jatropha + Pearl millet	27.76	24.71	20.49	18.76	23.90	22.72	9.14	8.94	26.82	25.34	68.39	64.87	28.61	28.64
C. D. at 5%	NS	0.15	NS	0.35	NS	0.15	NS	0.07	NS	0.67	NS	0.54	NS	NS

The data in parenthesis indicate % reduction over control.

TABLE 5
Growth yield attributes and seed yield of *Jatropha* during 2005-06 and 2006-07

Growth/yield characters	2004-05	2005-06		2006-07	
	Sole <i>Jatropha</i>	Sole <i>Jatropha</i>	Crops+ <i>Jatropha</i>	Sole <i>Jatropha</i>	Crops+ <i>Jatropha</i>
Plant height (m)	1.9	2.6	2.6	2.5	2.6
Primary branches/plant	7.9	30.6	29.2	36.5	34.2
Clusters/plant	12.2	24.9	20.4	13.8	14.6
Fruits/cluster	9.6	3.7	3.5	24.6	23.4
Seed yield(q/ha)	1.4	0.17	0.15	-	-

due to the fact that the *Jatropha* plants energy was mainly diverted towards vegetative growth which is evident from nearly fourfold increase in branches/plant after pruning in February, 2005. Only the fruits which set in early flowering (September) could mature before the frost. During the second year (2006-07) heading back of all the dead branches of *Jatropha* affected by frost resulted in emergence of a number of new branches resulting in excessive vegetative growth thereby delaying the onset of flowering to the month of October. Whereas, under normal conditions flowering starts by the end of August or early September. Due to delayed flowering, fertilization at low temperature in November and December was poor resulting in very shrivelled kernals with practically negligible oil content. Therefore, seeds were not picked from the plants to avoid wastage of manpower.

ACKNOWLEDGEMENTS

The authors are thankful to Dr. S. S. Bisla and D.P.S. Nandal, Department of Forestry, CCS HAU, Hisar for providing valuable guidance and suggestions during experimentation.

REFERENCES

- Batish, R. Daizy, R. K. Kohli, and H. P. Singh, (2011) : *J. Plant Biol.* **38** : 33-42.
- Divya, M. P., R. Santhi, and K. R. Ramesh, 2006 : *Indian J. Agroforestry.* **8** : 1-4.
- Joker, D and J. Jepsen, 2003 : *Jatropha curcas* L. seed leaflet No. 83 August 2003. Danida Forest Seed Centre, Denmark.
- Makkar, H. P. S., K. Becker, and B. Schmart, 2001 : Edible provenances of *Jatropha curcas* from Quintna Roo State of Mexico and effect of roasting on antinutrient and toxic factors in seeds. Institute for Animal Production in the Tropics and Subtropics (430), University of Hohenheim, D-70593 stuttgart, Germany.
- Rizvi, S. J. H., M. Tahir, V., Rizvi, R. K. Kohli, and A. Ansari, 1999 : *Critical Reviews Plant Sciences.* **18** : 773-796.
- Sharma, B. M. 2003 : *Adv. Arid. Legumes Res.* 279-284
- Singh, V.R.R; Dinesh Kumar; Sanjay Singh; Rites Sharma; Harish Chandra and Kshity Malhotra 2009 : *Indian Forester.* **135** : 287-289.