

## VARIABILITY OF SOIL INFILTRATION RATE IN DIFFERENT GRASS BASED CROPPING SYSTEM IN KYMORE PLATEAU SATPURA HILLS REGION

BHERU LAL KUMHAR\*, K. K. AGRAWAL, A. K. JHA AND H. K. RAI

College of Agriculture, Jabalpur  
Jawaharlal Nehru Agricultural University, Jabalpur-482004 (Madhya Pradesh), India  
\*(e-mail : [prajapatiagro09@gmail.com](mailto:prajapatiagro09@gmail.com))

(Received : 5 December 2021; Accepted : 30 December 2021)

### SUMMARY

A field study was conducted during two consecutive years (2018-19 and 2019-2000) at Research Farm, Department of Agronomy, Jawaharlal Nehru Krishi Vishwa Vidyalaya, Jabalpur to evaluate the infiltration behaviour of soil under eight grass based cropping system combinations. The recommended dose of 150 kg N, 80 kg P<sub>2</sub>O<sub>5</sub> and 40 kg K<sub>2</sub>O/ha was applied through urea, single super phosphate and muriate of potash. Results revealed that highest (6.1 and 11.6 m/hr) infiltration rate at the initiation (July, 2018) and end (June, 2020) of the study, respectively, was recorded under pearl millet napier hybrid in paired rows + ricebean - egyptian clover system, however, sole cropping of guinea grass (GG) system recorded lowest (5.0 and 7.4 mm/hr) infiltration rate at above respective times which were higher than that recorded before initiation of the study (4.5 mm/hr) during 2016.

**Key words :** Cumulative infiltration, grasses and cropping system, infiltration rate and vertisols

Infiltration is the hydrological process which specifies the knowledge of movement of water into the soil from surface sources such as precipitation and irrigation. Infiltration is the process of water entering in soil from the soil surface (Hillel, 1988) and the rate by which it enters into the soil is called the infiltration rate (Haghighi *et al.*, 2010). It is influenced by many internal and external factors, such as rainfall characteristics, slope gradients, soil properties, suction head, humidity, water content, types of impurities and surface sealing and cropping system which, makes infiltration hard to quantify (Herrada *et al.*, 2014). The infiltration rate of soil is reliant on several factors such as soil type, soil density, soil texture and moisture content (Angelaki *et al.*, 2013). Infiltration rate of soil is inversely proportion to the water-holding capacity of soil Singh *et al.*, 2014. Physical changes in soil caused by differential rooting behaviour of crops also affect the infiltration rate (Gupta and Gupta, 2008). Different cropping system has variable potentials to alter the soil properties (carbon content, aggregation, porosity, bulk density and water retention and release behaviors) and also affect the water infiltration rate of the soil. Overall the infiltration rate is higher at the beginning which decreases in elapsed time as it approaches the saturated hydraulic conductivity. The constant rate at which water enters into the soil after attaining saturation

is termed basic infiltration rate (Singh *et al.*, 2019). The intake rate relies upon the physical, chemical, and organic properties of the surface soil, the underlying dispersion of water in soil preceding irrigation, the transmission and redistribution of water over the surface. The measure of infiltration of water into the soil is an important indication concerning the efficiency of irrigation and drainage, optimizing the availability of water for plants, improving the yield of crops and minimizing erosion. The information about water transmission rate into the ground is vital concerning the productivity of water system and seepage, advancing the accessibility of water for the plants, enhancing the yield of harvests, limiting degradation of soil and wastage of the water. Therefore present study entitled “Variability of soil infiltration rate in different grass based cropping system in Kymore Plateau Satpura Hills region” was under taken.

### MATERIALS AND METHODS

Field study was conducted at Research Farm, Department of Agronomy, Jawaharlal Nehru Krishi Vishwa Vidyalaya, Jabalpur, Madhya Pradesh during 2018-2019 to 2019- 2020 to assess the variability in water infiltration rate of a Vertisol as affected by different grass based cropping system.

### Treatment details

The study was comprised of eight treatments of grass based cropping systems *i.e.* pearl millet napier (PN) hybrid sole crop ( $T_1$ ), guinea grass sole crop ( $T_2$ ), PN hybrid in paired rows + ricebean (*kharif*) - egyptian clover (*rabi*) ( $T_3$ ), PN hybrid in paired rows + *Desmanthus* ( $T_4$ ), PN hybrid in paired rows + *Sesbania grandiflora* ( $T_5$ ), guinea grass in paired rows + ricebean (*kharif*) - egyptian clover (*rabi*) ( $T_6$ ), guinea grass in paired rows + *Desmanthus* ( $T_7$ ) and guinea grass in paired rows + *Sesbania grandiflora* ( $T_8$ ) replicated thrice in randomized block design. The recommended dose of 150 kg N, 80 kg  $P_2O_5$  and 40 kg  $K_2O$ /ha was applied through urea, single super phosphate and muriate of potash. Study was started 01.06.2018 as pre-establish crop from 01.06.2016 by using three months old root slips for planting of both PN hybrid and guinea grass. All the observations were recorded and common package of practices was adopted for raising the crops.

### Measurement of infiltration rate

In this study, the infiltration rate was measured using double ring infiltrometer (Perroux and White, 1988) which consists of two concentric metal rings and gauge. The diameter of the inward and outward rings was 30 and 60 cm individually and both had an equivalent height of 30 cm. Double rings were used to eliminate the problems of over estimating the infiltration rate due to three dimensional flows. Rings are put concentrically and hammered with the help of rammer to ensure 10 cm insert of rings into the soil consistently and measuring gauge has been placed on the rings. A thin sheet of plastic was used inside the ring to avoid the surface soil disturbance due to splashing action of poured water. The water poured over the plastic sheet up to a depth of 15–20 cm. The




Fig. 1. Effect of grass based cropping systems on cumulative infiltration of a Vertisol during 2018-19.

plastic sheet was removed slowly and the outer ring filled with water up to the same level as inside or more but not less than inside water level. Infiltration readings were recorded using hook gauge fixed over the gauging stand and rate of fall in water level was measured at 30 minutes time interval till constant value of infiltration at two time intervals achieved (Pandey and Pandey, 2019). The readings of infiltration were plotted on a graph against the time to compute the infiltration rate.

### Data analysis

Analysis of data was carried out for every 30 minutes elapsed time. Test of significance was carried out by prepared ANOVA Table. Analysis of infiltration rate under eight grass based cropping system was also done.

## RESULTS AND DISCUSSION

### Cumulative infiltration (mm)

Data pertaining to variability in cumulative infiltration under different grass based cropping systems have been depicted in Fig. 2 and 3. Result indicated that highest cumulative infiltration (58.0 and 112.0 mm) at initiation (July 2018) and end (June 2020) of the study, respectively, was obtained under pearl millet napier hybrid in paired rows + ricebean - egyptian clover system and lowest (44.5 and 71.0 mm) at above respective time have been under sole cropping of guinea grass (GG) system.

### Infiltration rate (mm/h)

Data pertaining to variability of infiltration rates in different grass based cropping systems are given in Table 1 and depicted in Fig. 3. Result showed that both initial (July 2018) and final (June 2020) infiltration




Fig. 2. Effect of grass based cropping systems on cumulative infiltration of a Vertisol during 2019-2020.

TABLE 1  
Effect of grass based cropping systems on infiltration rate a Vertisol

Cropping Systems	Infiltration rate (mm/h)	
	2018-19	2019-20
T <sub>1</sub> –Pearl millet Napier (PN) hybrid sole cropping	5.04	7.50
T <sub>2</sub> –Guinea grass (GG) sole cropping	5.00	7.39
T <sub>3</sub> –PN hybrid in paired rows + Ricebean - Egyptian clover	6.08	11.63
T <sub>4</sub> –PN hybrid in paired rows + <i>Desmanthus</i>	5.92	9.52
T <sub>5</sub> –PN hybrid in paired rows + <i>Sesbania grandiflora</i>	5.44	8.16
T <sub>6</sub> –GG in paired rows + Ricebean - Egyptian clover	5.98	10.34
T <sub>7</sub> –GG in paired rows + <i>Desmanthus</i>	5.69	9.38
T <sub>8</sub> –GG in paired rows + <i>Sesbania grandiflora</i>	5.21	7.97
SEm ±	0.107	0.549
CD ( $p=0.05$ )	0.318	1.606
Initial value	4.5 mm/h	

rate was maximum (6.08 and 11.63 mm/h) under PN hybrid in paired rows + ricebean - egyptian clover cropping system which was statistically at par with guinea grass in paired rows + ricebean - egyptian clover system (5.98 and 10.34 mm/hr) and PN hybrid in paired rows + *Desmanthus* system (5.92 and 9.52 mm/hr) but significantly superior over those found under other cropping systems. Minimum infiltration rate (5.00 and 7.39 mm/h) was found under sole cropping of Guinea grass system which have been statistically on par with those obtained under PN hybrid sole cropping (5.04 and 7.50 m/h) and GG in paired rows + *Sesbania grandiflora* system (5.21 and 7.97 mm/h), respectively during both the years of experimentation.

The higher rate of infiltration under pearl millet napier hybrid in paired rows + ricebean - egyptian clover is may be due to deep, fibrous and tap root system, minimum tillage practices, surface covered throughout the year, stubble decomposition and timely irrigation. Increase in infiltration rate to enhanced soil aggregates, stable aggregates and content reduction in bulk density due to inclusion of legume crops in




Fig. 3. Effect of grass based cropping systems on infiltration rate of a Vertisol.

perennial grass based cropping system is directly related to soil infiltration (Srivastava *et al*, 2018). Reduction of bulk density enhance the soil porosity and volume leads to faster infiltration rate. Sand and clay content is the most crucial factor, which affects the soil infiltration rate. The bulk density (BD) and moisture content showed another possible input as these properties inversely related to infiltration characteristics (Pandey and Pandey, 2019). The organic carbon (OC) content of the soil showed a positive correlation with infiltration. The similar results confirmed by (Pandey and Pandey, 2018), (Verma and Sahu, 2007) and (Chouhan, *et al*, 2018). Higher rate of infiltration may be attributed to lower bulk density, higher organic carbon content and better soil aggregation which improve the movement of water in soil (Brar *et al*, 2015).

## CONCLUSIONS

On the basis forgoing results, it is concluded that in grass based cropping system, cumulative time is an essential parameter which affects the infiltration rate of the soil. The highest steady rate of infiltration was recorded under pearl millet napier hybrid in paired rows + ricebean - egyptian clover.

## ACKNOWLEDGEMENT

Authors greatly acknowledge, JNKVV Jabalpur, to provide all necessary facilities for conduction of research trial and EPCO, Bhopal for financial support as Chief Minister Fellowship on climate change.

## REFERENCES

- Angelaki, A., M. M. Sakellariou and C. Tzimopoulos, 2013 : Theoretical and experimental research of cumulative infiltration. *Transport in Porous Media*, **100**(2): 247-257.
- Brar, B. S., J. Singh, G. Singh and G. Kaur, 2015 : Effects of long term application of inorganic and organic fertilizers on soil organic carbon and physical properties in maize – wheat rotation. *Agronomy*, **5** : 220-238.
- Chouhan, B., H. K. Rai, A. Suryawanshi, S. Mandloi and R. Pandey, 2018 : Characterization of physical properties of Vertisols under long-term fertilizer experiment in soybean - wheat cropping system. *Journal of Pharmacognosy and Phytochemistry*, **7**(2): 596-601.
- Gupta, B. L. and A. Gupta, 2008 : Water resources systems and management, 2<sup>nd</sup> edn. Standard Publishers Distributors, Delhi, 510-535.
- Haghighi, F., M. Gorji, M. Shorafa, F. Sarmadian and M. H. Mohammadi, 2010 : Evaluation of some infiltration models and hydraulic parameters. *Spanish Journal of Agricultural Research*, **8**(1) : 210-217.
- Herrada, M. A., M. A. Gutiérrez and J. M. Montanero, 2014 : Modeling infiltration rates in a saturated/unsaturated soil under the free draining condition. *Journal of Hydrology*, **515** : 10-15.
- Hillel, D., 1998 : Environmental Soil Physics; Academic Press: New York, NY, USA, p120.
- Pandey, P. K. and V. Pandey, 2019 : Estimation of infiltration rate from readily available soil properties (RASPs) in fallow cultivated land. *Sustainable Water Resources Management*, **5** : 921-934.
- Perroux, K. and I. White, 1988 : Designs for disc permeameters. *Soil Science Society American Journal*, **52**(5): 1205-1215.
- Singh, B., P. Sihag and S. Dewal, 2019 : Modelling of the impact of water quality on the infiltration rate of the soil. *Applied Water Science*, **9**(15): 1-9.
- Singh, B., P. Sihag and D. Singh, 2014 : Study of infiltration characteristics of locally soils. *Journal of Civil Engineering Environment Technology*, **1** : 9-13.
- Srivastava, T. K., K. P. Singh, P. Singh, A. Suman, S. R. Singh, R. R. Verma, V. K. Singh and R. K. Singh, 2018 : Effect of bio-manures on soil quality, cane productivity and soil carbon sequestration under long-term sugarcane (*Saccharum officinarum*) plant - ratoon system in Indian subtropics. *Indian Journal of Agricultural Sciences*, **88**(11): 1696-1703.
- Verma, M. and M. L. Sahu, 2017 : Variability of soil infiltration under fifteen year old plantation in Jabalpur, Central India. *Indian Forester*, **143**(6) : 581-584.